

Find the radius (and interval) of convergence for

1. Recall the geometric series

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}, \quad |r| < 1$$

Let $a = 1$ and $r = x$

- (a) Write the sum of the series:

- (b) How would this be related to $f(x) = \frac{1}{1-x}$?

- (c) What is the interval of convergence?

- (d) Adapt the series so it is centered at -1

2. (a) Find a power Series for $f(x) = \frac{4}{x+2}$

Hint: write this in the form of $\frac{a}{1-r}$

- (b) What is the interval of convergence?

3. (a) Find a power series for $f(x) = \frac{1}{x}$ centered at 1

Hint: write this in the form of $\frac{a}{1-r}$

- (b) What is the interval of convergence?

4. (a) Find a power series for $f(x) = \frac{3x - 1}{x^2 - 1}$ centered at 0

Hint: Use partial fraction decomposition, then use the geometric series trick with both

- (b) What is the interval of convergence?

5. (a) Find a power series for $f(x) = \ln x$ centered at 1

Hint: Use integration! Didn't we do a series for $\frac{1}{x}$ before?

$$\ln x = \int \frac{1}{x} dx + C$$

- (b) What is the interval of convergence?

6. Find a power series for $f(x) = \arctan x$ centered at 0

(a) Recall

$$\frac{d}{dy} (\arctan y) = \frac{1}{1+y^2}$$

substitute $y = x^2$. Doesn't

$$f(x^2) = \frac{1}{1+x^2} = \sum_{n=0}^{\infty} (-1)^n x^{2n}?$$

Now integrate to find the series for $\arctan x$

(b) It can be shown that this series for $\arctan x$ converges for $x = \pm 1$. What is the series approximation for $\arctan 1$. Use your calc's sum(seq()) function to add a 100 or so terms. Is it close to $\frac{\pi}{4}$?

Answers:

3a. $\dots + \frac{x^8}{8} - \frac{x^6}{6} + \frac{x^4}{4} - \frac{x^2}{2} + 1$ Converges on the open interval $(-1, 1)$

3b. $\dots + \frac{x^8}{8} - \frac{x^6}{6} + \frac{x^4}{4} - \frac{x^2}{2} + 1$ Converges on the open interval $(0, 2)$

4a. $\dots + \frac{x^8}{8} - \frac{x^6}{6} + \frac{x^4}{4} - \frac{x^2}{2} + 1$ so $\frac{1}{1-x} + \frac{1}{1+x} = \frac{1-x}{1-x} + \frac{1+x}{1+x} = \frac{1-x+x}{1-x-x} = \frac{1}{1-x^2} = f(x)$

4b. $\dots + \frac{x^8}{8} - \frac{x^6}{6} + \frac{x^4}{4} - \frac{x^2}{2} + 1$ Converges on the open interval $(-1, 1)$

5a. To find $\arctan 1$, let $x = 1$ so $\arctan 1 = \ln = \frac{1}{1-x} + \frac{1}{1+x} = \frac{1}{1-x} + \frac{1}{1+x}$ so $\frac{1}{1-x} + \frac{1}{1+x} = \frac{1}{1-x^2} = f(x)$ Converges on the open interval $(0, 2)$ — see how integration can change the converge at the end points?

5b. $\dots + \frac{x^8}{8} - \frac{x^6}{6} + \frac{x^4}{4} - \frac{x^2}{2} + 1$ so $\arctan x = \frac{1}{1-x} + \frac{1}{1+x}$ so $\arctan x = \frac{1}{1-x} + \frac{1}{1+x}$ Converges on the open interval $(-1, 1)$

6b. Converges very slowly...