

Intro to Euler's Method (6.1b)

1. (2013 BC 5) Consider the differential equation $\frac{dy}{dx} = y^2(2x + 2)$. Let $y = f(x)$ be the particular solution to the differential equation with initial condition $f(0) = -1$

(a) Find $\lim_{x \rightarrow 0} \frac{f(x) + 1}{\sin x}$

- (b) Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $f\left(\frac{1}{2}\right)$.

- (c) Find $y = f(x)$, the particular solution to the differential equation with initial condition $f(0) = 1$.

2. Let $y = f(x)$ be the solutions to the differential equation $\frac{dy}{dx} = 2y - x$ with the initial condition $f(1) = 2$. What is the approximation for $f(0)$ obtained using Euler's method with two steps of equal length starting at $x = 1$?

- (a) $-\frac{5}{4}$
- (b) -1
- (c) $\frac{1}{4}$
- (d) $\frac{1}{2}$
- (e) $\frac{27}{4}$

3. Let $y = f(x)$ be the solutions to the differential equation $\frac{dy}{dx} = x - y - 1$ with the initial condition $f(1) = -2$. What is the approximation for $f(1.4)$ if Euler's method is used, starting at $x = 1$ with two steps of equal size?

- (a) -2
- (b) -1.24
- (c) -1.2
- (d) -0.64
- (e) 0.2

4. (2005 BC 4) Consider the differential equation $\frac{dy}{dx} = 2x - y$.

(a) On the axes provided, sketch a slope field for the given differential at the twelve points indicated, and sketch the solution curve that passes through the point $(0,1)$.

(b) The solution curve that passes through the point $(0,1)$ has a local minimum at $x = \ln(3/2)$. What is the y -value of this local minimum?

(c) Let $y = f(x)$ be the particular solution to the given differential equation with the initial condition $f(0) = 1$. Use Euler's method, starting at $x = 0$ with two steps of equal size, to approximate $f(-0.4)$. Show the work that leads to your answer.

(d) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Determine whether the approximation found in part (c) is less than or greater than $f(-0.4)$. Explain your reasoning.

