# Summary of Convergence Tests

NAME	STATEMENT	COMMENTS
Test for Divergence	If $\lim_{n \to \infty} a_n \neq 0$ , then $\sum a_n$ diverges.	If $\lim_{n \to \infty} a_n = 0$ , then $\sum a_n$ may or
	$\mu \rightarrow \infty$	may not converge.
P-Series Test	Let $\sum \frac{1}{n^p}$ be a series with positive terms, then (a) Series converges if $p>1$ (b) Series diverges if $p\leq 1$	This test can be used in conjunction with the comparison test for any $a_n$ whose denominator is raised to the $n^{th}$ power.
Direct Comparison Test	Let $\sum a_n$ and $\sum b_n$ be series with non-negative terms such that $a_1 \le b_1, a_2 \le b_2, a_3 \le b_3,$ if $\sum b_n$ converges, then $\sum a_n$ converges, and if $\sum a_n$ diverges, then $\sum b_n$ diverges.	This test works best for series whose formulas look very similar to the format needed for another test. Best example: when $a_n$ is a rational funct. WARNING: must use a second test to determine whether the built series converges or diverges.
Limit Comparison Test	Let $\sum a_n$ and $\sum b_n$ be series with positive terms such that $c = \lim_{n \to \infty} \frac{a_n}{b_n}$ if $0 < c < \infty$ , then both series converge, or both series diverge.	This is easier to apply than the Direct Comparison Test and works in the same type of cases. WARNING: must use a second test to determine whether the built series converges or diverges.
Integral Test	Let $\sum a_n$ be a series with positive terms, and let $f(x)$ be the function that results when <i>n</i> is replaced by <i>x</i> in the n <sup>th</sup> term of the associated sequence. If $f(x)$ is decreasing and continuous for $x \ge 1$ , then $\sum_{n=1}^{\infty} a_n$ and $\int_1^{\infty} f(x) dx$ both converge or both diverge.	This test only works for series that have positive terms. Try this test when $f(x)$ is easy to integrate.
Ratio Test	Let $\sum a_n$ be a series and suppose $\ell = \lim_{n \to \infty} \frac{a_{n+1}}{a_n}$ (a) Converges if $-1 < \ell < 1$ (b) Diverges if $\ell > 1$ or $\ell < -1$ (c) Test fails if $\ell = \pm 1$	Try this test when $a_n$ involves factorials or combinations of different types of functions.
Ratio Test for Absolute Convergence	Let $\sum a_n$ be a series and suppose $\ell = \lim_{n \to \infty} \left  \frac{a_{n+1}}{a_n} \right $ (a) Series converges if $\ell < 1$ (b) Series diverges if $\ell > 1$ (c) Test fails if $\ell = 1$	This is the default test because it is one of the easiest tests and it rarely fails. NOTE: the series need not have only positive terms nor does it have to be alternating.
Root Test	Let $\sum a_n$ be a series and suppose $\ell = \lim_{n \to \infty} \sqrt[n]{ a_n }$ (a) Series converges if $\ell < 1$ (b) Series diverges if $\ell > 1$ (c) Test fails if $\ell = 1$	This test is the most accurate, but not the easiest to use in many situations. Use this test when $a_n$ has n <sup>th</sup> powers.
Alternating Series Test	If $a_n > 0$ for all <i>n</i> , then the series $a_1 - a_2 + a_3 - a_4 + \dots$ or $-a_1 + a_2 - a_3 + a_4 - \dots$ Converges if: (a) $a_1 > a_2 > a_3 > a_4 > \dots$ (b) $\lim_{n \to \infty} a_n = 0$	This test <u>only</u> applies to alternating series. NOTE: $ S - S_n  \le  a_{n+1} $ which means that <b>error</b> of the partial sum of the first number of terms of the series is less than the absolute value of next term.

## Summary of Tests for Series

/ y \$	Test	Series	Converges	Diverges	Comment
	Test for Divergence	$\sum_{n=1}^{\infty} a_n$		$\lim_{n\to\infty}a_n\neq 0$	This test cannot be used to show convergence.
	Geometric Series	$\sum_{n=0}^{\infty} ar^n$	<i>r</i>   < 1	$ r  \ge 1$	Sum: $S = \frac{a}{1-r}$
	Telescoping Series	$\sum_{n=1}^{\infty} (b_n - b_{n+1})$	$\lim_{n\to\infty}b_n=L$		Sum: $S = b_1 - L$
a de la companya de l	p-Series	$\sum_{n=1}^{\infty} \frac{1}{n^p}$	<i>p</i> > 1	$p \leq 1$	
	Alternating Series	$\sum_{n=1}^{\infty} (-1)^{n-1} a_n$	$0 < a_{n+1} \le a_n$ and $\lim_{n \to \infty} a_n = 0$		Remainder: $ R_N  \leq a_{N+1}$
	Integral (f is continuous, positive, and decreasing)	$\sum_{n=1}^{\infty} a_n,$ $a_n = f(n) \ge 0$	$\int_{1}^{\infty} f(x)  dx \text{ converges}$	$\int_{1}^{\infty} f(x)  dx \text{ diverges}$	Remainder: $0 < R_N < \int_N^\infty f(x) dx$
	Root	$\sum_{n=1}^{\infty} a_n$	$\lim_{n\to\infty}\sqrt[n]{ a_n } < 1$	$\lim_{n\to\infty}\sqrt[n]{ a_n } > 1$	Test is inconclusive if $\lim_{n \to \infty} \sqrt[n]{ a_n } = 1.$
	Ratio	$\sum_{n=1}^{\infty} a_n$	$\lim_{n \to \infty} \left  \frac{a_{n+1}}{a_n} \right  < 1$	$\lim_{n \to \infty} \left  \frac{a_{n+1}}{a_n} \right  > 1$	Test is inconclusive if $\lim_{n \to \infty} \left  \frac{a_{n+1}}{a_n} \right  = 1.$
	Direct Comparison $(a_n, b_n > 0)$	$\sum_{n=1}^{\infty} a_n$	$0 < a_n \le b_n$ and $\sum_{n=1}^{\infty} b_n$ converges	$0 < b_n \le a_n$ and $\sum_{n=1}^{\infty} b_n$ diverges	
	$\begin{array}{l} \text{Limit Comparison} \\ (a_n, b_n > 0) \end{array}$	$\sum_{n=1}^{\infty} a_n$	$\lim_{n \to \infty} \frac{a_n}{b_n} = L > 0$ and $\sum_{n=1}^{\infty} b_n$ converges	$\lim_{n \to \infty} \frac{a_n}{b_n} = L > 0$ and $\sum_{n=1}^{\infty} b_n$ diverges	

#### Strategies for Testing Series

You have now studied ten tests for determining the convergence or divergence of an infinite series. (See the summary in the table on page 593.) Skill in choosing and applying the various tests will come only with practice. Below is a useful checklist for choosing an appropriate test.

Guidelines for Testing a Series for Convergence or Divergence

- 1. Does the *n*th term approach 0? If not, the series diverges.
- 2. Is the series one of the special types—geometric, *p*-series, telescoping, or alternating?
- 3. Can the Integral Test, the Root Test, or the Ratio Test be applied?
- 4. Can the series be compared favorably to one of the special types?

In some instances, more than one test is applicable. However, your objective should be to learn to choose the most efficient test.

### **EXAMPLE 5** Applying the Strategies for Testing Series

Determine the convergence or divergence of each series.

a.	$\sum_{n=1}^{\infty} \frac{n+1}{3n+1}$	<b>b.</b> $\sum_{n=1}^{\infty} \left(\frac{\pi}{6}\right)^n$	<b>c.</b> $\sum_{n=1}^{\infty} n e^{-n^2}$	<b>d.</b> $\sum_{n=1}^{\infty} \frac{1}{3n+1}$
e.	$\sum_{n=1}^{\infty}  (-1)^n  \frac{3}{4n  +  1}$	<b>f.</b> $\sum_{n=1}^{\infty} \frac{n!}{10^n}$	$\mathbf{g} \cdot \sum_{n=1}^{\infty} \left( \frac{n+1}{2n+1} \right)$	$\left(\frac{1}{1}\right)^n$

#### Solution

- a. For this series, the limit of the nth term is not 0 (a<sub>n</sub>→<sup>1</sup>/<sub>3</sub> as n→∞). Thus, by the nth-Term Test, the series diverges.
- **b.** This series is geometric. Moreover, because the common ratio of the terms is less than 1 in absolute value  $(r = \pi/6)$ , you can conclude that the series converges.
- **c.** Because the function  $f(x) = xe^{-x^2}$  is easily integrated, you can use the Integral Test to conclude that the series converges.
- **d.** The *n*th term of this series can be compared to the *n*th term of the harmonic series. After using the Limit Comparison Test, you can conclude that the series diverges
- **e.** This is an alternating series whose *n*th term approaches 0. Because  $a_{n+1} \le a_n$ , you can use the Alternating Series Test to conclude that the series converges.
- f. The *n*th term of this series involves a factorial, which indicates that the Ratio Test may work well. After applying the Ratio Test, you can conclude that the series diverges.
- **g.** The *n*th term of this series involves a variable that is raised to the *n*th power, which indicates that the Root Test may work well. After applying the Root Test, you can conclude that the series converges.